

ANTI-FOAM

CONCENTRATED ANTI-FOAMING AGENT

Batch #:

Date of
Manufacture:

DO NOT STORE AAC ANTI-FOAM MIXED WITH WATER
OR OTHER CHEMICALS

MIX ENOUGH AAC ANTI-FOAM FOR 1 DAY'S APPLICATION.

AAC ANTI-FOAM is compatible with a range of commonly used spray chemicals. When mixing with other chemicals, always mix a small quantity and check for precipitate (fall out) in mixture. If precipitation occurs, DO NOT proceed with application.

CONDITIONS OF SALE

To the extent permitted by law, all conditions and warranties and statutory or other rights of action which buyer or any other user may have against AUSTRALIAN AGRICULTURAL CHEMICALS are hereby excluded. AUSTRALIAN AGRICULTURAL CHEMICALS hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information provided by AUSTRALIAN AGRICULTURAL CHEMICALS or on its behalf unless it is established that such information or advice was provided negligently and that the product has been used strictly as directed. AUSTRALIAN AGRICULTURAL CHEMICALS's liability shall in all circumstances be limited to replacement of product or a refund of the purchase price thereof.

NOTE

The suggested application rates are designed for typical Australian conditions and act as a guide only. Differences in soil types, climatic conditions, water quality, application methods and processes and therefore necessitate corrections to ensure optimum results. Best practice requires that applications under extreme weather conditions such as temperatures over 30°C, high humidity, frost, rain should be avoided. It's recommended that prior to applying to a crop or area for the first time, or in combination with other chemicals, a small test area should be sprayed and observed prior to the total crop spray. It is recommended that leaf (sap) tests are conducted on a regular basis to monitor actual plant nutrient availability during each growing cycle. Soil tests at least once per year are essential.

DIRECTIONS FOR USE:

General Use Rate: 20 - 50 mL / 100 L of spray solution.
Apply to tank mix in situations where excessive foaming occurs in mixing and application.

STORAGE & DISPOSAL

Store in original container, tightly closed in a safe place. Wash out and destroy empty containers.

PROTECTION OR WILDLIFE, FISH & ENVIRONMENT

Do not contaminate streams, rivers or waterways with mixes or used containers.

SAFETY DIRECTIONS

Mild irritant. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a Doctor or local poisons information centre. Additional information is listed in the Material Safety Data Sheet.

CONTENTS:

5 L 10 L 20 L 200 L 1000 L

**AUSTRALIAN
AAC
AGRICULTURAL
CHEMICALS**

1/3972 Pacific Hwy Loganholme
Queensland, Australia. 4129

Phone: +61 7 3806 5733
Fax: +61 7 3806 5733

www.aussie-ag.com.au