



POSTERITY – the fungicide that holds strong for many moons.




syngenta®

Introducing POSTERITY fungicide, the latest SDHI in the turf market that provides exceptional Dollar Spot control. It is a great option to supercharge your tank mix and deal with Dollar Spot, with power that lasts. POSTERITY sets a new standard, holding stronger for many moons.

What POSTERITY offers:

- Long lasting curative and preventative control against difficult-to-control, Dollar Spot.
- A formulation that is compatible with a wide range of commonly used products including PRIMO MAXX so you can supercharge your tank mix.
- Balanced distribution within the plant and outstanding rainfastness – it binds quickly and strongly to the plant's wax layer and then slowly penetrates into the plant tissue.
- A step change in Dollar Spot control vs traditional controls such as DMI fungicides containing for example, Propiconazole and Tebuconazole.
- Smarter chemistry that is exempt from poison scheduling.
- A highly effective option at rates as low as 500 mL/ha to 1 L/ha.

POSTERITY: A step change in Dollar Spot Control



Dedicate® is a Registered Trademark of the Bayer Group. Trial Reference: PBICobbity_ABN_2016_WG

To achieve great results, accurate application is important

Target	Syngenta Nozzle	Optimal Water Rate	Optimal Pressure Range
Leaf and Crown F	XC04	400 L/ha	2-4 bar

Dollar Spot Control Program

To combat **Dollar Spot**, start the season strong with an early application of POSTERITY when conditions are favourable for disease outbreak or as an early curative application at first signs of disease. Favourable conditions include: temperatures between 15°C and 30°C, continuous high humidity, cool nights and intense dews.

For **resistance management** and broader-spectrum control, Syngenta agronomic programs strategically rotate POSTERITY with DAPONIL WEATHERSTIK, BANNER MAXX and other trusted products to help prevent Dollar Spot.

For **continued protection** from late Summer through Autumn, turn to POSTERITY for curative and extended preventative applications. Truly the next generation carboxamide (SDHI), a powerfully simple choice.



UNTREATED



POSTERITY 0.5 L/ha (2 applications)

Trial ref: Clark2016_RU

Disease	Rate	Critical Comments
Dollar Spot <i>(Clariireedia homoeocarpa)</i> (formerly <i>Sclerotinia homoeocarpa</i>)	0.5 L/ha	Preventative treatment: apply prior to symptom development, when conditions are favourable to disease. Apply at 14 day intervals.
	0.75 - 1 L/ha	Early curative / preventative treatment: Apply prior to symptom development or at first signs of disease. Re-apply after 28 days. Apply no more than 2 sequential applications.

Refer to the registered label for full details and instructions for use.

F = Foliar applied **So** = Soil applied **Su** = Surfactant required

Active Ingredient
POSTERITY contains 200 g/L pydiflumetofen, a member of the carboxamide (SDHI) fungicide group
GROUP 7 FUNGICIDE

Formula & Packaging
500 mL pack size
Suspension concentrate (SC) that mixes completely with water and may be tank mixed with many commonly used fungicides, insecticide, herbicides and fertilisers.



For more information, ask your Syngenta Agent or call Syngenta Customer Service on 1800 022 035 or visit www.greencast.com.au